

U-391

ADDRESS		KJELLER N-2007 Kjeller, Norway	HALDEN N-1751 Halden, Norway	AVAILABILITY Private confidential
TELEPHONE		(02)712560 - 713560	(031)83100	
TELEX		76361 energ n	76335 energ n	
REPORT TYPE	REPORT NO.	IFE/KRF-84/112		DATE 1984-11-06
	REPORT TITLE	RESERVOIR GAS GEOCHEMISTRY ON SAMPLE FROM WELL 31/5-2		DATE OF LAST REV.
	CLIENT	STATOIL, NORSK HYDRO A/S, SAGA A/S		REV. NO.
	CLIENT REF.			NUMBER OF PAGES
SUMMARY				DISTRIBUTION
<p>A rather heavy <math>\delta^{13}\text{C}</math> propane value together with the uncommon relationships between the <math>\delta^{13}\text{C}</math> values of propane and the butanes can in our view only be explained by effects from bacterial degradation in the reservoir.</p> <p>This in accordance with the conclusion given in the reports on well 31/3-1, 31/5-1/31/5-2, 31/6-2, 31/6-5 and in IFE report No. IFE/KR/F-84/096.</p> <p>This biodegradation was followed by a period of renewed supply of predominantly dry gas. A crossplot between the <math>\delta^{13}\text{C}</math> methane and the <math>\delta\text{D}</math> methane values indicates that this superimposed gas is condensate associated. The hydrogen of the methane is isotopically lighter than reported earlier for the other Troll field wells. This may suggest that the gas is supplied from a source with different isotopic composition of the kerogens.</p>				Oppdragsgivere, 10 Andresen, B. Berg, J.O. Brevik, E.M. Gaudernack, B. Garder, K. Råheim, A.
BA 84-2477-1 19 NOV 1984 <b>REGISTRERT</b> <b>OLJEDIREKTORATET</b>				SUMMARY Aamodt, N-G. Project leaders Halden Sectionheads Kjeller Library
KEYWORDS				
	NAME		DATE	SIGNATURE
PREPARED BY	Einar M. Brevik Björg Andresen Arne Råheim		1984-11-06	<i>Einar Brevik Björg Andresen Arne Råheim</i>
REVIEWED BY				
APPROVED BY	Karen Garder		1984-11-06	<i>Karen Garder</i>

ANALYTICAL PROCEDURE

The sample was received in a pressurized gas bottle. The natural gas has been separated into the different gas components by a Carlo-Erba 4200 instrument. The hydrocarbon gas components were oxydized in separate CuO-ovens, which enables us to collect several times when the concentration of a gas component is low. The combustion products CO<sub>2</sub> and H<sub>2</sub>O were frozen into collection vessles and separated. The isotopic measurements were performed on a Finnigan Mat 251 mass spectrometer. Our δ<sup>13</sup>C-value on NBS-22 is -29.77 +/- .06 ‰.

RESULTS

The stable isotope results of the natural gas from well 31/5-2 are given in the following table :

Table 1. Isotopic composition of gas samples from well 31/5-2.

	<sup>C<sub>1</sub></sup> δ <sup>13</sup> C    δD	C <sub>2</sub>	C <sub>3</sub>	iC <sub>4</sub>	nC <sub>4</sub>	CO <sub>2</sub> δ <sup>13</sup> C    δ <sup>18</sup> O
DST I	-47.5    -255	- 26.6	- 16.0	- 21.4	- 21.2	+4.8    -13.2

The composition in % of the gas sample is given in table 2. The result has not been normalized to 100%, the rest is air.

1577-1581

30/8-84

Table 2. Composition of gas from well 31/5-2 % of C<sub>1</sub>-C<sub>4</sub>

	Sample 1
C <sub>1</sub>	78.0
C <sub>2</sub>	4.2
C <sub>3</sub>	0.5
iC <sub>4</sub>	0.4
nC <sub>4</sub>	0.05
CO <sub>2</sub>	0.9

#### INTERPRETATION

The rather heavy  $\delta^{13}\text{C}$  propane values together with the uncommon relationships between the  $\delta^{13}\text{C}$  values of propane and the butanes can in our view only be explained by effects from bacterial degradation in the reservoir.

A new, relatively dry gas has migrated into the reservoir after the bacterial degradation had stopped. The new gas mixed with the gas left after the bacterial degradation.

This makes it impossible to use Alan T. James' method \* to indicate the maturity of the source.

It is, however, likely that the methane component of the natural gas is representative for the new, relatively dry gas. This assumption is based on our experience on bacterial degradation from laboratory experiments. The carbon and hydrogen isotopic composition of the gas can therefore be plotted in a  $\delta^{13}\text{C}$  methane vs.  $\delta\text{D}$  methane plot (Schoell, 1983)\*\*, figure 1. This indicates that the superimposed gas is condensate associated and possibly formed at high maturity in the condensate field of the oil window. The hydrogen of the methane is isotopically lighter than reported earlier for the other Troll field wells. This may be due to mixtures of gases from different origins or this part of the Troll field may be supplied by gas from a source or sources with different isotopic composition of the kerogens.

\* James, Alan T. (1983) : Correlation of Natural Gas by Use of Carbon Isotopic Distribution Between Hydrocarbon Components, A.A.P.G. Vol. 67, No. 7, July, 1983.

\*\* Schoell, M. 1983 : Genetic Characertization of Natural Gases. A.A.P.G. December, 1983.




Figure 1 a. Variations of molecular composition in natural gases related to the isotope variations of methane.




Figure 1 b. Carbon and hydrogen isotope variation in methane.

The principle for the genetic characterization of natural gases is that the primary gases (B-biogenic gas, T-associated gas, TT-non-associated gas) are defined by fields of compositional variations. These primary gases may become mixed and form various mixtures "M" of intermediate composition. "TT(m)" and "TT(h)" are non associated gases from marine source rocks and coal gases from N.W. Germany respectively, compositional shifts due to migration are indicated by arrows Md (deep migration) and Ms (shallow migration) respectively. "T" are gases associated with petroleum in an initial phase of formation. "T<sub>c</sub>" are gases associated with condensates. (Schoell, 1983).