



## CONTINENTAL SHELF INSTITUTE

Håkon Magnussons gt. 1B — N-7000 Trondheim — Telephone (075) 15660 — Telex 55548

REPORT TITLE/ TITTEL Analyses of cores from well 6407/1-2			
CLIENT/ OPPDRAGSGIVER Statoil			
RESPONSIBLE SCIENTIST/ PROSJEKTANSVÄRLIG Magnar Björny			
AUTHORS/ FORFATTERE Magnar Björny, Peter B. Hall and Torun Vinge			
DATE/ DATO 1.3.83	REPORT NO./ RAPPORT NR. 05.0089	NO. OF PAGES/ ANT. SIDER 53	NO. OF ENCLOSURES/ ANT. BILAG

RECEIVED  
CONTINENTAL SHELF INSTITUTE

SUMMARY/ SAMMENDRAG

Ten core samples from well 6407/1-2 were analysed. All the samples contained a light crude oil.

KEY WORDS/ STIKKORD


## CONTENTS

	Page
EXPERIMENTAL AND DESCRIPTION OF INTERPRETATION LEVELS	1
RESULTS AND DISCUSSIONS	4
Rock-Eval Pyrolysis	4
Thermal Evaporation with Capillary GC Analysis, and Pyrolysis GC	4
Table 1: Rock-Eval Pyrolyses	5
Extraction and Chromatographic Separation	6
Saturated Hydrocarbons	7
Aromatic Hydrocarbons	7
Table 2: Concentration of Extractable Organic Matter and Chromatographic Fractions	8
Table 3: Concentration of Extractable Organic Matter and Chromatographic Fractions (mg/g TOC)	9
Table 4: Composition in % of Material Extracted from the Rock	10
Table 5: Lithology and Total Organic Carbon Measurements	11
CONCLUSIONS	13
Figures:	
Thermal Evaporation Chromatograms	
Pyrolysis Gas Chromatograms	
Saturated Hydrocarbon Chromatograms	
Aromatic Hydrocarbon Chromatograms	

## EXPERIMENTAL

### Total Organic Carbon (TOC)

Samples were crushed in a centrifugal mill. Aliquots of the samples were then weighed into Leco crucibles and treated with hot 2N HCl to remove carbonate and washed twice with distilled water to remove traces of HCl. The crucibles were then placed in a vacuum oven at 50°C and evacuated to 20 mm Hg for 12 hrs. The samples were then analysed on a Leco E C 12 carbon analyser, to determine the total organic carbon (TOC).

The results are shown in table 5 with the lithological description.

### Rock-Eval Pyrolysis

100 mg crushed sample was put into a platinum crucible whose bottom and cover are made of sintered steel and analysed on a Rock-Eval pyrolyser. The results are shown in table 1.

### Pyrolysis Gas Chromatography (Py-GC)

#### Thermoextraction.

20-30 mg of whole rock sample was placed in a boat shaped sample probe and thermoextracted in a stream of helium at 300°C for 5 minutes.

#### Py-GC

20-30 mg of thermoextracted whole rock sample was programmed pyrolysed in helium (260°C to 520°C at 35°C/min.) in a furnace type pyrolyzer. The outlet of the pyrolyzer was directly connected to a splitter (30:1) and a fused silica capillary column. The pyrolysis product was trapped in a cooled (liq. Nitrogen) U-shaped section at the front of the column.

The outlet of the splitter was directly connected to a FID detector and the course of the pyrolysis could be followed by the detector response of the bulk pyrolysis product (30:1) which was recorded as a

broad peak. At the end of the pyrolysis the pyrolysis product was injected on to the capillary column at ambient temperature (by removing the nitrogen bath) and analysed under the GC conditions given below.

#### GC-conditions

Column: 25m OV-1, I.D. 0.3 mm, fused silica capillary column.

Carrier gas: Helium with inlet pressure 10 psi. Flow; ca. 1.5. ml/min.

Oven programme: 40<sup>o</sup> - 270<sup>o</sup>C at 4<sup>o</sup>C/min.

#### Extractable Organic Matter (EOM)

From the TOC results samples were selected for extraction. Of the selected samples, approximately 100 gm of each was extracted in a flow through system (Radke et al., 1978, Anal. Chem. 49, 663-665) for 10 min. using dichloromethane (DCM) as solvent. The DCM used as solvent was distilled in an all glass apparatus to remove contaminants.

Activated copper filings were used to remove any free sulphur from the samples.

After extraction, the solvent was removed on a Buchi Rotavapor and transferred to a 50 ml flask. The rest of the solvent was then removed and the amount of extractable organic matter (EOM) determined.

#### Chromatographic Separation

The extractable organic matter (EOM) was separated into saturated fraction, aromatic fraction and non hydrocarbon fraction using a MPLC system with hexane as eluant (Radke et al., Anal. Chem., 1980). The various fractions were evaporated on a Buchi Rotavapor and transferred to glassvials and dried in a stream of nitrogen. The various results are given in tables 2-4.

Gas Chromatographic Analyses

The saturated and aromatic hydrocarbon fractions were each diluted with n-hexane and analysed on a HP 5730 A gas chromatograph, fitted with a 25 m OV101 glass capillary column and an automatic injection system. Hydrogen (0.7 ml/min.) was used as carrier gas and the injection was performed in the split mode (1:20).

## RESULTS AND DISCUSSION

### Rock-Eval Pyrolysis

Quick screening analysis of the ten core samples was performed by pyrolysing aliquots of the samples on a Rock-Eval II instrument. Since these are reservoir samples, ordinary parameters from Rock-Eval pyrolysis such as hydrogen and oxygen indices are meaningless, and we will discuss the non normalised values. The results are shown in Table 1.

The Rock-Eval data shows a slightly decreasing trend with increasing depth down to 3671.9m. The samples between 3676.5 and 3780.5m show very variable values. This could be due to the porosity/permeability of the samples analysed. All of the samples contain free hydrocarbons. None of the samples have a large  $S_2$  peak, indicating that if there are any kerogen or asphaltenes in the samples, these will only give a minor contribution.

### Thermal Evaporation with Capillary GC Analysis, and Pyrolysis GC

This method is not a quantitative method, but qualitative. It gives a gas chromatographic picture of the free hydrocarbons in the sample released during heating (at  $300^{\circ}\text{C}-S_1$ ) and the hydrocarbons produced by pyrolysis of the kerogen, if there are any ( $S_2$ ).

The samples in this study can be placed in two major categories (A and B) and one intermediate group (category C). Category A includes those with a large unresolved envelope in the high molecular weight area, and an abundance of  $nC_{26}-nC_{30}$  n-alkanes, and category B comprises those samples which do not have this unusual feature.

Category A; Samples from 3661.4m, 3664.85m, 3676.5m, 3683.65m, 3688.5m and 3692.45m belong in this group. These have a good abundance of  $C_{10}-C_{25}$  n-alkanes, indicating a relatively light oil. The pristane/ $nC_{17}$  ratio is high while the pristane/phytane ratio is low when compared to a large proportion of oils found in the North Sea. The loss of n-alkanes below  $nC_{10}$  is due to the permeability of the cores, the light hydrocar-

TABLE 1.

ROCK EVAL PYROLYSES

IKU No.	DEPTH m/ft	S1	S2	S3	TOC (%)	HYDR. INDEX	OXYGEN INDEX	OIL OF GAS CONTENT S1+S2	PROD. INDEX S1+S2	TEMP. MAX (C)
A 3129	3661.40	2.83	0.18	0.11	0.06	300	183	3.01	0.94	413
A 3130	3664.85	2.44	0.18	0.12	0.24	69	46	2.62	0.93	407
A 3131	3671.90	2.15	0.27	0.08	0.23	117	35	2.42	0.89	412
A 3132	3676.50	3.12	0.24	0.13	0.25	96	52	3.36	0.93	407
A 3133	3680.10	1.88	0.18	0.09	0.18	100	50	2.06	0.91	410
A 3134	3683.65	0.92	0.29	0.03	0.20	145	15	1.21	0.76	411
A 3135	3688.50	4.06	0.37	0.15	0.10	370	150	4.43	0.92	399
A 3136	3692.45	2.02	0.27	0.10	0.17	159	59	2.29	0.88	415
A 3137	3696.90	2.63	0.39	0.08	0.24	162	33	3.02	0.87	412
A 3138	3700.00	2.09	0.24	0.12	0.24	100	50	2.33	0.90	405

DATE : 15 - 2 - 83.

bons evaporate when the core is brought up to the rig. The gas chromatograms show a large unresolved envelope and an increase in the n-alkane distribution above  $nC_{25}$ . This could be due to some residues such as asphaltenes etc. which start to crack and give off hydrocarbons at  $300^{\circ}C$ .

Category B; The two most typical samples for this category, which have a smooth n-alkane distribution and no unresolved envelope are 3671.9m and 3700m. The front part of the gas chromatograms is similar to that found for the samples from category A, while there is no unresolved envelope and no increase in the n-alkane distribution above  $nC_{25}$ . This indicates that these samples do not contain any material which decomposes at  $300^{\circ}C$ , producing hydrocarbons.

Category C; (Intermediate between A and B). Two samples, 3680.10m and 3696.90m have similar front-end distributions to the other samples but with an increase in n-alkanes from  $nC_{20}$  and without the unresolved envelope. These two samples have a smaller pristane/phytane ratio than the other samples. The sample from 3680.10m has a far stronger front-end distribution than the rest of the samples.

The pyrolysis gc patterns of the organic remains vary only slightly for the ten core samples. All the samples have a small methane peak and small amounts of hydrocarbons up to  $nC_{25}$  together with an unresolved envelope and large single peaks in the  $nC_{25}$ - $nC_{30}$  range, (this is similar to the distribution found for category A, in the thermal extraction analysis). The amounts vary from sample to sample, but the feature is quite distinguishable in all samples.

The samples from category B, (as defined by the thermal evaporation) also have large unresolved envelopes for the pyrolysis gc's, indicating that this phenomena is present in all the samples although it is not seen in some when thermal evaporation is performed.

#### Extraction and Chromatographic Separation

Extraction of the samples indicates that down to 3676.5m there is a general increase of extractable hydrocarbons with increasing depth. The samples from 3680.1m and 3683.65m have a low abundance of extract-



able hydrocarbons while the sample from 3688.5m has the highest abundance of all the analysed samples. The extractability decreases for the sample from 3692.45m compared with the sample above, then to return to increasing slightly with increasing depth. The abundance of extractable organic material follows the same trends as the abundance of extractable hydrocarbons. This data indicates that the reservoir down to approximately 3676.5m has a high abundance of extractable organic matter with a large percentage of hydrocarbons, probably reflecting good porosity/permeability for the section. The samples from 3680.10-3683.65m have a far lower extractability. This could be due to lighter, less permeable sediment. The samples just below have far higher extractability and higher percentage of hydrocarbons than any other samples indicating a build up of hydrocarbon just below the "seal". The lower part of the analysed section (3692.45-3700m) is similar to the upper part. The extraction results are shown in tables 2-4.

#### Saturated Hydrocarbons

The gas chromatograms of the saturated hydrocarbon fractions vary only slightly for the analysed samples. All the gc's show a front-biased, smooth n-alkane distribution with moderate pristane/nC<sub>17</sub> ratios and higher pristane/phytane ratios. The pristane/phytane ratios are higher than those normally found in the North Sea.

#### Aromatic Hydrocarbons

The aromatic hydrocarbon distribution in these samples is dominated by C<sub>1</sub>-naphthalenes (A) and C<sub>2</sub>-naphthalenes (B). Other prominent compounds include Naphthalene (N) C<sub>3</sub>-naphthalenes (C), phenanthrene P, C,- and C<sub>2</sub>-phenanthrene (D and E). The peaks marked with an asterisk mark the location of organic sulphur compounds. This distribution is typical for light-medium gravity oils probably with no alteration from biodegradation or water washing (probably derived from type II or mixed type II/III kerogen).

T A B L E : 2.

CONCENTRATION OF EOM AND CHROMATOGRAPHIC FRACTIONS

I	I	I	I	I	I	I	I	I	I	I
I	IKU-No	DEPTH	Rock	Extr.	EOM	Sat.	Aro.	HC	Non	TOC
I		(m)	(g)	(mg)	(mg)	(mg)	(mg)	(mg)	(mg)	(%)
I	A 3129	3661.40	53.5	248.6	128.5	45.3	173.8	74.8	0.23	
I	A 3130	3664.85	50.6	230.4	116.8	27.1	143.9	86.5	0.27	
I	A 3131	3671.90	52.8	233.0	116.6	66.0	182.6	50.4	0.26	
I	A 3132	3676.50	35.0	196.1	88.8	47.2	136.0	60.1	0.23	
I	A 3133	3680.10	50.9	145.3	77.4	15.5	92.9	52.4	0.09	
I	A 3134	3683.65	43.7	71.0	36.0	18.4	54.4	16.6	0.33	
I	A 3135	3688.50	53.3	361.8	220.2	108.0	328.2	33.6	0.24	
I	A 3136	3692.45	54.8	259.5	120.4	59.5	179.9	79.6	0.23	
I	A 3137	3696.90	52.1	226.5	119.8	59.3	179.1	47.4	0.24	
Y	A 3138	3700.00	49.4	315.8	136.3	68.8	205.1	110.7	0.21	

DATE : 15 - 2 - 83.

T A B L E : 3.

CONCENTRATION OF EOM AND CHROMATOGRAPHIC FRACTIONS

(mg/g TOC)

I	I	I	I	I	I	I	I	I	I	I
I	IKU-No	DEPTH	EOM	Sat.	Ar.o.	HC	Non	HC	I	I
I		(m)							I	I
I	A 3129	3661.40	2018.8	1043.5	367.9	1411.4	607.4	I	I	I
I	A 3130	3664.85	1687.1	855.3	198.4	1053.7	633.4	I	I	I
I	A 3131	3671.90	1697.9	849.7	481.0	1330.6	367.3	I	I	I
I	A 3132	3676.50	2436.0	1103.1	586.3	1689.4	746.6	I	I	I
I	A 3133	3680.10	3169.9	1688.6	338.2	2026.7	1143.2	I	I	I
I	A 3134	3683.65	492.1	249.5	127.5	377.1	115.1	I	I	I
I	A 3135	3688.50	2825.7	1719.8	843.5	2563.3	262.4	I	I	I
I	A 3136	3692.45	2058.9	955.3	472.1	1427.3	631.5	I	I	I
I	A 3137	3696.90	1811.4	958.1	474.2	1432.3	379.1	I	I	I
I	A 3138	3700.00	3046.0	1314.7	663.6	1978.3	1067.7	I	I	I

DATE : 15 - 2 - 83.

T A B L E : 4.

COMPOSITION IN % OF MATERIAL EXTRACTED FROM THE ROCK

IKU-No	DEPTH (m)	Sat EOM	ArO EOM	HC EOM	SAT ArO	Non HC EOM	HC Non HC
A 3129	3661.40	51.7	18.2	69.9	283.7	30.1	232.4
A 3130	3664.85	50.7	11.8	62.5	431.0	37.5	166.4
A 3131	3671.90	50.0	28.3	78.4	176.7	21.6	362.3
A 3132	3676.50	45.3	24.1	69.4	188.1	30.6	226.3
A 3133	3680.10	53.3	10.7	63.9	499.4	36.1	177.3
A 3134	3683.65	50.7	25.9	76.6	195.7	23.4	327.7
A 3135	3688.50	60.9	29.9	90.7	203.9	9.3	976.8
A 3136	3692.45	46.4	22.9	69.3	202.4	30.7	226.0
A 3137	3696.90	52.9	26.2	79.1	202.0	20.9	377.8
A 3138	3700.00	43.2	21.8	64.9	198.1	35.1	185.3

DATE : 15 - 2 - 83.



# Lithology and Total Organic Carbon measurements

TABLE NO.: 5.  
WELL NO.: 6407/1-2

Sample	Depth (m)	TOC	Lithology
A-3129	3661.4	0.23	Sandstone, light brown, fine-medium grained quartz, angular mostly white some brown, micaceous, a residual black bitumen, which is non-fluor - very slight white cut. General (not overall) yellow (bright) - white fluorescence, with overall immediate fair white cut (not streaming). Hard, non calcareous kaolinite?, cemented - silica.
A-3130	3664.85	0.27	Sandstone, as above, more medium grained with micaceous shale laminae (carbonaceous) very pale yellow-white, immediate intense cut, streaming in part, more intense than A-3129.
A-3131	3671.90	0.26	Sandstone, as above, micaceous laminae, extremely hard, less intense cut than A-3130, no shale laminae.
A-3132	3676.50	0.23	Sandstone, light brown, fine-medium grained, hard, general bright yellow fluor, non-micaceous, immediate intense white cut. Quartz cement, little bitumen, clean sand with subangular grains to sub rounded in part.
A-3133	3680.10	0.09	Sandstone, darker brown some mica laminae again similar to A-3129, silty in part, distinct oil staining, cut as A-3132.




# Lithology and Total Organic Carbon measurements

TABLE NO.: 5.  
WELL NO.: 6407/1-2

Sample	Depth (m)	TOC	Lithology
A-3134	3683.65	0.33	Sandstone, darker brown as above thick mica laminae, cut as A-3132.
A-3135	3688.50	0.24	Sandstone, medium grained - some coarse quartz, little mica, lighter brown, slightly more brittle, oil stained clearly, fluor bright yellow, immediate intense white cut.
A-3136	3692.45	0.23	Fine-medium grained generally as above, (A-3135) white cut/fluor intens yellow.
A-3147	3696.90	0.24	Medium-coarse grained, hard, otherwise as above. Yellow fluor is darker yellow hue. There is less cut.
A-3138	3700.00	0.21	As above medium grained, fairly brittle, immediate intense (streaming in part) white cut.

Distribution of EOM and EHC with depth.




CONCLUSIONS

All ten samples from well 6407/1-2 contained free hydrocarbons, indicating that the samples are from a reservoir with light oil. Variation in abundance of hydrocarbons indicates a strong variation in the permeability of the analysed samples.


Large unresolved envelopes and large abundances of  $nC_{20}-nC_{30}$  indicate that there might be significant amounts of asphaltenes/residual oil which decompose at  $300^{\circ}C$  or higher.




Thermal Evaporation  
A - 3129  
Core 1  
3661.40m




Thermal Evaporation  
A - 3130  
Core 1  
3664.85m



Thermal Evaporation  
A - 3131  
Core 1  
3671.90m



Thermal Evaporation  
A - 3132  
Core 1  
3676.50m




Thermal Evaporation


A - 3133

Core 2


3680.10m




Thermal Evaporation  
A - 3134  
Core 2  
3683.65m



Thermal Evaporation  
A - 3135  
Core 2  
3688.50m




Thermal Evaporation  
A - 3136  
Core 3  
3692.45 m







Thermal Evaporation  
A - 3137  
Core 3  
3696.90m




Thermal Evaporation  
A - 3138  
Core 3  
3700.00m




Pyrolysis GC  
A - 3129  
Core 1  
3661.40m




Pyrolysis GC  
A - 3130  
Core 1  
3664.85m




Pyrolysis GC  
A - 3131  
Core 1  
3671.90m




Pyrolysis GC  
A - 3132  
Core 1  
3676.50m



Pyrolysis GC  
A - 3133  
Core 2  
3680.10m




Pyrolysis GC  
A - 3134  
Core 2  
3683.65m







Pyrolysis GC  
A - 3135  
Core 2  
3688.50m




Pyrolysis GC  
A - 3136  
Core 3  
3692.45m




Pyrolysis GC  
A - 3137  
Core 3  
3696.90m




Analysis : 0089A3129S1 Sample #: 1 Injection #: 1  
Sample Name : R-3129,S,6407/1-2,TV Maximum value : 5116




Analysis : 050089S1    Sample #: 1    Injection #: 1  
Sample Name : R-3130,S,6407/1-2,TY    Maximum value : 4447




Analysis : 0089R3131S1 Sample #: 1 Injection #: 1  
Sample Name : R-3131, S, 6407/1-2, TV Maximum value : 16383



Analysis : 0089R3132S1 Sample f: 1 Injection f: 1  
Sample Name : R-3132,S,6407/1-2,TV Maximum value : 5468




Analysis : 0089R3133S1 Sample #: 1 Injection #: 1  
Sample Name : R-3133,S,6407/1-2,TV Maximum value : 4528






Analysis : 0089R3134S1 Sample #: 1 Injection #: 1  
Sample Name : R-3134,S,6407/2-1,TV Maximum value : 4454




Analysis : 0089R3135S1 Sample #: 1 Injection #: 1  
Sample Name : A-3135,S,6407/1-2,TV Maximum value : 8027




Saturated HC  
A - 3135  
3688.50m

Analysis : 0089A3136S1 Sample #: 1 Injection #: 1  
Sample Name : R-3136,S,6407/1-2,TV Maximum value : 5640




Saturated HC  
A - 3136  
3692.45m

Analysis : 0089A3137S1 Sample #: 1 Injection #: 1  
Sample Name : R-3137,S,6407/1-2,TV Maximum value : 13938




Analysis : 0089R3138S1    Sample #: 1    Injection #: 1  
Sample Name : A-3138,S,6407/1-2,TV    Maximum value : 5137




Saturated HC  
A - 3138  
3700.00m

Analysis : 0089A3129A1 Sample #: 1 Injection #: 1  
Sample Name : A-3129,A,6407/1-2,TV Maximum value : 16383




Aromatic HC  
A - 3129  
3661.40m

Analysis : 0089A3130R1 Sample #: 1 Injection #: 1  
Sample Name : R-3130, A, 6407/1-2, TV Maximum value : 11358




Analysis : 0089R3131R1 Sample #: 1 Injection #: 1  
Sample Name : R-3131, R, 6407/1-2, TV Maximum value : 16383







Analysis : 0089R3132R1 Sample #: 1 Injection #: 1  
Sample Name : R-3132, R, 6407/1-2, TV Maximum value : 6602




Analysis : 0089R3133R2 Sample #: 1 Injection #: 1  
Sample Name : R-3133, R, 6407/1-2, TV Maximum value : 9513




Analysis : 0089R3134R1 Sample f: 1 Injection f: 1  
Sample Name : R-3134, R, 6407/1-2, TV Maximum value : 294




Analysis : 0089R3135R1 Sample #: 1 Injection #: 1  
Sample Name : A-3135, R, 6407/1-2, TV Maximum value : 605



Analysis : 0089R3136R1 Sample #: 1 Injection #: 1  
Sample Name : A-3136, R, 6407/1-2, TV Maximum value : 5055



Analysis : 0089A3137A1 Sample #: 1 Injection #: 1  
Sample Name : R-3137,A,6407/1-2,TV Maximum signal (%): 18.14



Aromatic HC  
A - 3137  
3696.90m

Analysis : 0089A3138A1 Sample #: 1 Injection #: 1  
Sample Name : R-3138, R, 6407/1-2, TV Maximum value : 2810

