Encl. 2 Lithological and biostratigraphical data sheet

WELL: NOR 31/2-4 INTERVAL: 1320.0-1663.9M CORES: 1-28 DRILLERS DEPTH

SCALE: 1:200

LITHOLOGICAL LEGEND

SEDIMENTARY BEDDING

- ≈ UNBEDDED, MASSIVE
- ≡ WELL BEDDED
- = FAIRLY WELL BEDDED
- (=) SLIGHTLY BEDDED
- ≈ IRREGULAR, WAVY BEDDING
- ∠ X-BEDDING
- ∠ X-BEDDING LOW ANGLE
- FLASER BEDDING
- LENTICULAR BEDDING
- GRADED BEDDING

SEDIMENTARY STRUCTURES

- ≤ DISTURBED/FAULTED
- SLUMPED, CONTORTED
- LOAD CASTS
- ~ ASYMMETRICAL RIPPLES
- PEBBLE LINEATION

BIOLOGIC CHARACTERISTICS

- CHURNED, BIOTURBATED
- BURROWS
- BORED SURFACE
- SHELL,FOSSIL LINEATION
- UNSPEC. BROKEN FOSSILS,ANGULAR
- GASTROPODS
- LAMELLIBRANCHS(PELECYPODS,BIVALVES)
- BELEMNITES
- ☆ CRINOIDS
- FORAMINIFERA,GENERAL
- PLANT REMAINS
- PLANT FRAGMENT LINEATION

ACCESSORY MINERALS

GC	GLAUCONITE
MC	MICA
PY	PYRITE
SD	SIDERITE

RELATIVE ABUNDANCE INDICATOR

SCARCE
COMMON
ABUNDANT

→ UNLABELLED SEALED SAMPLES INDICATED ON (GECO) CORE PHOTOGRAPHS

NB IN THE LITHOLOGY CALCAREOUS SANDSTONE IS DENOTED AS SANDY LIMESTONE FOR REASONS OF CLARITY

AUTHORS: LEVELL-WILLIAMSON & DE WEERD (KSEPL), WHITAKER & NOORDERMEER (SIPM), LARSEN (NORSKE SHELL)

PALYNOLOGICAL QUALIFIERS

SOM STRUCTURELESS ORGANIC MATTER

* % OF REMAINING SUB-CATEGORIES ARE NOT DIFFERENTIATED INTO SMALL, MEDIUM OR LARGE

† VOL. OF ORGANIC RESIDUE-FILL IN MEASURED AMOUNT CALCULATED PER 10 GM

DIVERSITY

V. LOW = L
MEDIUM = M
V. HIGH= H

NO. OF DOMINANTS

1, 2, 3 ETC

DEGRADATION

POOR = P
MEDIUM = M
GOOD = G
CARBONIZED = B

ROUNDING

V. COMMON = C
MEDIUM = M
RARE = R

BROKEN

V. COMMON = C
MEDIUM = M
RARE = R

PALAEONTOLOGICAL LEGEND

SPECIMEN NO

■ = >100 SPECIMENS
● = 21-100 SPECIMENS
○ = 6-20 SPECIMENS
/ = 2-5 SPECIMENS
• = SINGLE SPECIMEN

NF = NO FORAMINIFERA

Koninklijke/Shell
Exploratie en Produktie Laboratorium

LITHOLOGICAL AND BIOSTRATIGRAPHICAL DATA (DD)
WELL 31/2-4
NORSKE SHELL

SCALE 1:200

INV. 929227		Date: NOVEMBER '82
Rep: RKER 82.258	Encl: 2	Draw.no: 36212